

Name:



Date of Birth (MM/DD/YYYY):

2021-2022 UCLA EXTENSION STUDY ABROAD IMMUNIZATION REQUIREMENTS

This form is required for all international students pursuing Study Abroad in UCLA and UCLA Extension credit and certificate programs. Please complete and return this form at least 30 days before the program start date.

LAST	FIRST	
REQUIRED IMMUNIZATIONS		
Tdap Vaccine • Tetanus/Diphtheria WITH Pertussis (whooping cough)	ONE DOSE ON OR AFTER SEVEN YEARS OF AGE FOR NON-HEALTH CARE PROFESSIONAL STUDENTS ONE DOSE IN THE LAST 10 YEARS required for health care professional students Dose Date: (Please note: The requirement is Tdap and not Td or Dtap.)	
• Measles, Mumps & Rubella	YOU MUST HAVE TWO DOSES WITH THE FIRST DOSE BEING ON OR AFTER YOUR FIRST BIRTHDAY. Dose 1 Date: (must be on or after your first birthday) (Dose 1 & 2 must be AT LEAST 28 days apart) Dose 2 Date: If unable to obtain proof of vaccination, you must obtain a blood titer test. * ATTACH A COPY OF YOUR LAB REPORT POSITIVE Measles IgG Antibody Titer Titer Date: POSITIVE Mumps IgG Antibody Titer Titer Date: POSITIVE Rubella IgG Antibody Titer Titer Date:	
Varicella (Chicken Pox) Vaccine	YOU MUST HAVE 2 DOSES WITH THE FIRST DOSE BEING ON OR AFTER YOUR FIRST BIRTHDAY. Dose 1 Date: (must be on or after your first birthday) (Dose 1 & 2 must be AT LEAST 28 days apart) Dose 2 Date: IF YOU HAD THE DISEASE AS A CHILD OR IF YOU ARE UNABLE TO OBTAIN PROOF OF VACCINATION, YOU MUST OBTAIN A BLOOD TITER TEST. POSITIVE Varicella IgG Antibody Titer Titer Date: • If you have a negative or indeterminate titer, obtain one dose of varicella vaccine and repeat titer four weeks later. If titer is still negative, receive a second dose of varicella vaccine.	
Meningococcal Vaccine • MCV4 (Menactra or Menveo brand preferred) REQUIRED for ALL students 21 yrs or younger	THE MOST RECENT DOSE MUST BE ON OR AFTER THE 16TH BIRTHDAY. Dose 1 Date: Dose 2 Date:	



Tuberculosis (TB) Questionnaire

		Date	of Birth (MM/DD/YYYY):
LAST	FIRST		
Please answer the following questions: O Have you ever had a positive TB skin or blood test? O Have you ever had close contact with anyone who was sick with TB? O Are you from or have you ever lived or traveled in one of the following are: Mexico, South or Central America, Eastern Europe, Asia, the Middle East, or			
If all questions are answered NO , you have completed your TB Assessment.			
If any questions are answered YES , then you must also have your health treatment for TB or negative TB test results. This must be completed and program. If TB results are required, the TB test must be taken no more	d submitted to UCLA Extensi	ion thirty (30) da	
Tuberculosis (TB) Assessment This part of the form more RISK FACTORS: (please ask student and check any that apply) 1. Immunosuppressed (HIV/AIDS), organ transplant, or on immunos 2. History of abnormal chest x-ray suggestive of TB disease 3. Does the student have signs or symptoms of active tuberculosis of (Cough more than 3 weeks, chest pain, unexplained weight loss, for the student have signs or symptoms.	suppressant medication Ye Ye disease? Ye		th care provider.
If no, proceed to 4 or 5. If yes, proceed with additional evaluation to exsputum evaluation as indicated, and show results below. 4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine,	6. Chest X-Ray (required	l if TST or IGRA is p	positive)
sputum evaluation as indicated, and show results below. 4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no	6. Chest X-Ray (required	l if TST or IGRA is p	positive)
4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is immunosuppressed. 10 mm induration is considered positive if coming		d if TST or IGRA is p	positive)
4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is	6. Chest X-Ray (required ○ Date of chest x-ray: ○ Result: normal If chest x-ray is a	d if TST or IGRA is p m abnormal bnormal, please s	oositive) onth/ day/ year (including scars, and old
sputum evaluation as indicated, and show results below. 4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is immunosuppressed. 10 mm induration is considered positive if coming from a high-risk area or has other high-risk conditions (IV drug use, chronic renal disease, cancer, diabetes, malabsorption or GI bypass).	6. Chest X-Ray (required ○ Date of chest x-ray: ○ Result: normal If chest x-ray is a Sputum Results (AFE	abnormal bnormal, please s	onth/ day/ year (including scars, and old granulomatous changes) submit the following results.
4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is immunosuppressed. 10 mm induration is considered positive if coming from a high-risk area or has other high-risk conditions (IV drug use, chronic renal disease, cancer, diabetes, malabsorption or GI bypass). O Date TST test was given: month/day/year	6. Chest X-Ray (required Date of chest x-ray: Result: normal If chest x-ray is a Sputum Results (AFE #1 Date	abnormal monormal monormal monormal monormal monormal monormal.	positive) onth/ day/ year (including scars, and old granulomatous changes) submit the following results. equired if chest x-ray abnorm _Culture_
sputum evaluation as indicated, and show results below. 4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is immunosuppressed. 10 mm induration is considered positive if coming from a high-risk area or has other high-risk conditions (IV drug use, chronic renal disease, cancer, diabetes, malabsorption or GI bypass). O Date TST test was given: month/day/year O Date TST test was read: month/day/year	6. Chest X-Ray (required Date of chest x-ray: Result: normal If chest x-ray is a Sputum Results (AFE #1 Date #2 Date	abnormal mand bnormal, please sand culture x 3 rangers	positive) onth/ day/ year (including scars, and old granulomatous changes) submit the following results. equired if chest x-ray abnorm _Culture_
4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is immunosuppressed. 10 mm induration is considered positive if coming from a high-risk area or has other high-risk conditions (IV drug use, chronic renal disease, cancer, diabetes, malabsorption or GI bypass). O Date TST test was given: month/day/year O Result: mm induration	6. Chest X-Ray (required Date of chest x-ray: Result: normal If chest x-ray is a Sputum Results (AFE #1 Date #2 Date	bnormal, please s AFB AFB AFB	positive) ponth/ day/ year (including scars, and old granulomatous changes) submit the following results. equired if chest x-ray abnorm Culture Culture Culture
4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is immunosuppressed. 10 mm induration is considered positive if coming from a high-risk area or has other high-risk conditions (IV drug use, chronic renal disease, cancer, diabetes, malabsorption or GI bypass). O Date TST test was given: month/day/year O Result: mminduration O Interpretation: negative positive	6. Chest X-Ray (required Date of chest x-ray: Result: normal If chest x-ray is a Sputum Results (AFE #1 Date #2 Date #3 Date 7. Treatment for Latent	abnormal meases and culture x 3 reaches AFB AFB	positive) ponth/ day/ year (including scars, and old granulomatous changes) submit the following results. equired if chest x-ray abnorm Culture Culture Culture
4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is immunosuppressed. 10 mm induration is considered positive if coming from a high-risk area or has other high-risk conditions (IV drug use, chronic renal disease, cancer, diabetes, malabsorption or GI bypass). O Date TST test was given: month/day/year O Result: mminduration O Interpretation: negative positive BBlood Test (Interferon Gamma Release Assay-IGRA) The TB	6. Chest X-Ray (required Date of chest x-ray: Result: normal If chest x-ray is a Sputum Results (AFE #1 Date #2 Date #3 Date 7. Treatment for Latent O Medication(s)	abnormal manabnormal, please sand culture x 3 range AFB AFB AFB	positive) ponth/ day/ year (including scars, and old granulomatous changes) submit the following results. equired if chest x-ray abnorm CultureCultureCulture
4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is immunosuppressed. 10 mm induration is considered positive if coming from a high-risk area or has other high-risk conditions (IV drug use, chronic renal disease, cancer, diabetes, malabsorption or GI bypass). O Date TST test was given: month/day/year O Result: mminduration O Interpretation: negative positive	6. Chest X-Ray (required Date of chest x-ray: Result: normal If chest x-ray is a Sputum Results (AFE #1 Date #2 Date #3 Date 7. Treatment for Latent	abnormal manabnormal, please sand culture x 3 range AFB AFB AFB	onth/ day/ year (including scars, and old granulomatous changes) submit the following results. equired if chest x-ray abnorm _CultureCultureCultureCulture
4. Tuberculin Skin Test (TST) If there is no history of BCG Vaccine, TST results should be recorded as millimeters (mm) of induration. If no induration, write "0." Five mm is considered positive if there is a history of abnormal chest x-ray, recent exposure to active TB disease, or is immunosuppressed. 10 mm induration is considered positive if coming from a high-risk area or has other high-risk conditions (IV drug use, chronic renal disease, cancer, diabetes, malabsorption or GI bypass). O Date TST test was given: month/day/year O Date TST test was read: month/day/year O Result: positive mm induration B Blood Test (Interferon Gamma Release Assay-IGRA) The TB od test may be done instead of TST. Strongly recommended if there	6. Chest X-Ray (required Date of chest x-ray: Result: normal If chest x-ray is a Sputum Results (AFE #1 Date #2 Date #3 Date 7. Treatment for Latent O Medication(s)	abnormal meases and culture x 3 r AFB AFB TB (if applicab	positive) ponth/ day/ year (including scars, and old granulomatous changes) submit the following results. equired if chest x-ray abnorm CultureCultureCultureCulture

	*NOTE: These vaccinations are recommended BUT NOT required to be compliant with enrollment.		
 HPV Vaccine Human Papilloma	RECOMMENDED FOR ALL STUDENTS (ALL GENDERS) UP TO THE AGE OF 26 HPV 4 Dose 1 Date: Dose 2 Date: Dose 3 Date:	RECOMMENDED FOR ALL STUDENTS (ALL GENDERS) UP TO THE AGE OF 26 HPV 9 Dose 1 Date: Dose 2 Date: Dose 3 Date:	
Meningococcal B Vaccine • Trumemba or Bexsero	RECOMMENDED FOR AGES 16-23 AFTER DISCUSSIO Dose 1 Date: Dose 2 Date: (Trumemba is either a two-dose or three-dose series Dose 3 Date:		
Hepatitis A Vaccine • 2 dose series	Dose 1 Date: (Dose 2 should be administered 6-12 months follow Dose 2 Date:	ring first dose.)	
Polio Vaccine • 4 dose series	Dose 1 Date: Dose 2 Date: Dose 3 Date: Dose 4 Date:		
Pneumococcal Vaccine PCV13 +/or PPSV23 based on health history	,	of asthma, diabetes, smokers and those with care provider	

UCLA Extension has no responsibility for verifying the accuracy of the information provided on this form.*

I ATTEST THAT ALL DATES AND IMMUNIZATIONS LISTED ON THIS FORM ARE CORRECT AND ACCURATE		
Student's Signature:	Date:	
Provider's Signature:	Practice Stamp:	
Provider's Name:(Physician/PA/NP/RN)	Date:	

176706-21 Page 3 of 3

^{*}As an educational institution, UCLA Extension treats the immunization information provided herein as personal confidential information in a student's record protected under the Family Educational Rights and Privacy Act of 1974 (FERPA). Thereby, this immunization information will be stored securely and not released to any outside entity.